



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

312. Proposed by J. A. CAPARO, C. E., Notre Dame University, Notre Dame, Ind.

Two roots of the cubic $x^3 - px^2 + qx - c = 0$ are equal. Find their value in terms of p , q , and c .

I. Solution by GEORGE W. HARTWELL, University of Kansas, Lawrence, Kas.; V. M. SPUNAR, M. and E. E., East Pittsburg, Pa.; and G. I. HOPKINS, Instructor in Mathematics and Astronomy, Manchester High School, N. H.

Let m , m , n be the three roots. Then

$$2m + n = p \dots (1); \quad 2mn + m^2 = q \dots (2); \quad m^2 n = c \dots (3).$$

Solving (1) and (2) for m and n we have,

$$m = \frac{p \pm \sqrt{(p^2 - 3q)}}{3}, \quad n = \frac{p \mp 2\sqrt{(p^2 - 3q)}}{3}.$$

Substituting these values in (3),

$$(2p^2 - 6q) \left(\frac{p \pm \sqrt{(p^2 - 3q)}}{3} \right) = pq - 9c \dots (4). \text{ But } \frac{p \pm \sqrt{(p^2 - 3q)}}{3} = m.$$

$$\text{Therefore, from (4), } m = \frac{pq - 9c}{2p^2 - 6q}.$$

II. Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.; J. SCHEFFER, A. M., Hagerstown, Md.; S. LEFSCHETZ, Wilkensburg, Pa.; and the PROPOSER.

$$\text{Let } f(x) = x^3 - px^2 + qx - c = 0, \quad f'(x) = 3x^2 - 2px + q = 0.$$

If $f(x)$ has two equal roots, $f'(x)$ contains one, and hence the greatest common divisor of $f(x)$ and $f'(x)$ gives one of the equal roots. Now if

$$(pq - 9c)(27c + 4p^3 - 15pq) = 4q(3q - p^2)^2, \text{ or} \\ 18cpq + p^2q^2 - 4cp^3 - 4q^3 - 27c^2 = 0,$$

then $2(3q - p^2)x + pq - 9c$ is the greatest common divisor of $f(x)$ and $f'(x)$.

Therefore, $x = (9c - pq) / [2(3q - p^2)]$ is one of the equal roots.


GEOMETRY.

339. Proposed by G. E. BROCKWAY, Boston, Mass.

Of all triangles that can be inscribed in a given triangle, that formed by joining the feet of the altitudes has the minimum perimeter. Prove by means of the straight line and circle.

I. Solution by J. SCHEFFER, A. M., Hagerstown, Md.

Assuming the points N and O in the sides AC and BC , $NM+MO$ is a minimum, if $\angle NMA = \angle OMB$; for letting fall the perpendicular NR and extending it to S by its own length, OMS becomes a straight line. It follows from this that in the case of MNO being a triangle of minimum perimeter, $\angle NMA = \angle BMO = \alpha$, $\angle ANM = \angle CNO = \beta$, $\angle MOB = \angle NOC = \gamma$.



Now, $\alpha + \beta = 180^\circ - A$, $\alpha + \gamma = 180^\circ - B$, $\beta + \gamma = 180^\circ - C$.
 $\therefore \alpha + \beta + \gamma = \frac{1}{2}(540^\circ - 180^\circ) = 180^\circ$. $\therefore \alpha = C$, $\beta = B$, $\gamma = A$.
 Consequently, the triangle MNO is the pedal triangle.

II. Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

When we wish to find a point P on a given line so that the sum of the distances $PR+PS$ to two given points, is a minimum, it is easy to show that PS and PR must make equal angles with the given line. Let M, L, K be the feet of the altitudes; M on AB , L on AC , K on BC . Then if M, L are fixed, K is the point on BC such that $KL+KM$ is a minimum, since KL, KM make equal angles with BC . Similarly, for M, K and L, K fixed in turn, respectively. $LM+LK$ is a minimum and $ML+MK$ is a minimum for the reason cited above.


This seems the easiest and simplest proof.

340. Proposed by J. H. MEYERS, S. J., Sacred Heart College, Augusta, Ga.

Given trapezoid $ABCD$. Prolong AB and CD , the non-parallel sides, to meet in E . On AE as diameter construct semi-circle $AHGE$. With BE as radius construct arc BG . Draw GK perpendicular to AE . Bisect AH at L . Erect KH perpendicular to AE . Construct arc HM with HE as radius. Draw MN perpendicular to DC . Prove that MN bisects the trapezoid $ABCD$, angles ADC and BCD being right angles.*

Solution by J. SCHEFFER, A. M., Hagerstown, Md.

We may generalize this theorem by drawing the line MN parallel to AD , instead of perpendicular to DC .



$$\triangle ADE : \triangle EMN = AE^2 : EM^2 = AE^2 : EH^2 = AE^2 : AE \times EL = AE : EL.$$

$$\therefore \triangle ADE - \triangle EMN : \triangle EMN = AE - EL, \text{ or}$$

$$AMND : \triangle EMN = AL : EL \dots (I).$$

$$\triangle EMN : \triangle EBC = ME^2 : BE^2 = KE^2 : GE^2 = AE \times EL : AE \times EK = EL : EK.$$

$$\therefore \triangle EMN - \triangle EBC : \triangle EMN = EL - EK : EL, \text{ or}$$

$$MBCN : \triangle EMN = LK : EL \dots (II).$$

$$\text{Comparing (I) and (II), } AMND : MBCN = AL : LK = 1 : 1.$$

*The reading of this problem has been slightly changed to correspond to the figure. ED. F.